



Mathematics Colloquium

Computer driven questions, pre-theorems and theorems in geometry

Moira Chas (Stony Brook University)

Abstract

Several numbers can be associated to a free homotopy class X of closed curves on a surface S with boundary and negative Euler characteristic. Among these,

- the self-intersection number of X (this is the smallest number of times a representative of the X crosses itself),
- the word length of X ,
- the length of the geodesic corresponding to X ,
- the number of free homotopy classes of a given word length the mapping class group orbit of X .

The interrelations of these numbers exhibit many patterns when explicitly determined or approximated by running a variety of algorithms in a computer.

We will discuss how these computations lead to counterexamples to existing conjectures and to the discovery of new patterns. Some of these new patterns, so intricate and unlikely that they are certainly true (even if not proven yet), are "pre-theorems". Many of these pre-theorems later became theorems. An example of such a theorem states that the distribution of the self-intersection of free homotopy classes of closed curves on a surface, appropriately normalized, sampling among given word length, approaches a Gaussian when the word length goes to infinity. An example of a counterexample (no pun intended!) is that there exists pairs of length equivalent free homotopy classes of curves on a surface S that have different self-intersection number. (Two free homotopy classes X and Y are length equivalent if for every hyperbolic metric M on S , $M(X) = M(Y)$).

Wednesday, 19 October 2016, 4:15pm

Smith Hall 204

Tea and refreshments will be served at 4pm.